# Neofitos Stefanides, M.D., P.C.

neostefanidesMD@yahoo.com

44-01 Francis Lewis Blvd L3B1205 Franklin Avenue Suite 15030-74 31st st. 2nd fl.Bayside, NY 11361Garden City, NY 11530Astoria, NY 11102Ph: 718.989.8515Ph: 718.989.8515Ph: 718.989.8515Fax: 718.626.0102Fax: 718.626.0102Fax: 718.626.0102

Name:	Date:
Diagnosis:	Date of Surgery:

## Rehabilitation Protocol: Arthroscopic SLAP Repair

### Phase I (Weeks 0-4)

Sling immobilization at all times except for showering and rehab under guidance of PT Range of Motion −AAROM → AROM as tolerated

- $\circ~$  Restrict motion to  $140^\circ$  of Forward Flexion,  $40^\circ$  of External Rotation and Internal Rotation to stomach
- $\circ\;$  No Internal Rotation up the back/No External Rotation behind the head Therapeutic Exercise
- o Wrist/Hand Range of Motion
- o Grip Strengthening
- o Isometric Abduction, Internal/External Rotation exercises with elbow at side
- No resisted Forward Flexion/Elbow Flexion (to avoid stressing the biceps origin)
  Heat/Ice before and after PT sessions

#### Phase II (Weeks 4-6)

Discontinue sling immobilization

Range of Motion – Increase Forward Flexion, Internal/External Rotation to full motion as tolerated Therapeutic Exercise

- o Advance isometrics from Phase I to use of a theraband within AROM limitations
- o Continue with Wrist/Hand Range of Motion and Grip Strengthening
- o Begin Prone Extensions and Scapular Stabilizing Exercises (traps/rhomboids/levator scapula)
- Gentle joint mobilization

Modalities per PT discretion

#### Phase III (Weeks 6-12)

Range of Motion - Progress to full AROM without discomfort

Therapeutic Exercise – Advance theraband exercises to light weights (1-5 lbs)

- o 8-12 repetitions/2-3 sets for Rotator Cuff, Deltoid and Scapular Stabilizers
- o Continue and progress with Phase II exercises
- o Begin UE ergometer

Modalities per PT discretion

# Neofitos Stefanides, M.D., P.C.

neostefanidesMD@yahoo.com

44-01 Francis Lewis Blvd L3B

Bayside, NY 11361 Ph: 718.989.8515 Fax: 718.626.0102 1205 Franklin Avenue Suite 150

Garden City, NY 11530 Ph: 718.989.8515 Fax: 718.626.0102 30-74 31st st. 2nd fl. Astoria, NY 11102 Ph: 718.989.8515 Fax: 718.626.0102

## Phase IV (Months 3-6)

Range of Motion – Full without discomfort

Therapeutic Exercise – Advance exercises in Phase III (strengthening 3x per week)

- o Sport/Work specific rehabilitation
- o Return to throwing at 4.5 months
- $\circ\;$  Return to sports at 6 months if approved

Modalities per PT discretion